

# Capitainer® B Vanadate

Product code: 210-0052 50 pcs box

The ultimate sampling solution for PEth analysis

- The risk of post sampling PEth formation eliminated.
- All the benefits of fingerblood and dried samples.
- Volumetric, solves the problem with traditional DBS.
- More reliable than venous blood samples.
- Easy patient centric sampling.



## The Capitainer® solution:

- Based on traditional DBS technology.
- Exact volume of blood through advanced microfluidic technology.
- Hematocrit independent.
- Eliminates the risk of overfilling.
- Colour indicator for successful sampling.
- Send by regular mail, the dried sample is non-biohazard.

<1,2 grams plastics per device.  
>80% of the card is paper based.



## Increased accuracy in PEth analysis

Capitainer®B Vanadate is the only volumetric microsampling device scientifically shown to enable accurate PEth measurement without post sampling formation. It is based on our proprietary technology\* integrating the phospholipase D (PLD) inhibitor sodium metavanadate ( $\text{NaVO}_3$ ) in the DBS paper disc. The inhibitor stops all formation of PEth post sampling [1]. This makes Capitainer®B Vanadate the ultimate solution for PEth monitoring, outperforming traditional venous samples.

\* patent pending PCT/EP2021/068386

### Specifications:

- Sample volume: 2 x 10  $\mu\text{l}$ .
- Precision, CV < 5%
- Size: 78 x 43 x 3 mm.
- Operating temperature range: +15°C to +35°C.
- Hematocrit range: 25% to 55%.
- Precut DBS disc, 96-well plate compatible



### About Phosphatidylethanol (PEth)

PEth is formed in the membrane of red blood cells in the presence of ethanol. The half-life of PEth is around 4 days in blood and it therefore accumulates with frequent alcohol consumption. This makes PEth a direct marker for long term alcohol use. However, PEth has been shown to form post sampling if phospholipase D (PLD) activity is not inhibited. Post sampling formation can result in inaccurate results.

To learn more  
contact us at  
[info@stratech.co.uk](mailto:info@stratech.co.uk)

Hope to hear  
from you!

[1] Beck, O., Mellring, M., Löwbeer, C. et al. Measurement of the alcohol biomarker phosphatidylethanol (PEth) in dried blood spots and venous blood—importance of inhibition of post-sampling formation from ethanol. Anal Bioanal Chem (2021). <https://doi.org/10.1007/s00216-021-03211-z>